



## Research Publication Repository

<http://publications.wehi.edu.au/search/SearchPublications>

Author's accepted manuscript version	Uren RT, Iyer S, Kluck RM. Pore formation by dimeric Bak and Bax: an unusual pore? <i>Philosophical Transactions of the Royal Society of London. Series B, Biological Sciences</i> . 2017 372(1726) pii: 20160218.
Final published version	DOI: 10.1098/rstb.2016.0218 <a href="http://rstb.royalsocietypublishing.org/content/372/1726/20160218.long">http://rstb.royalsocietypublishing.org/content/372/1726/20160218.long</a>
Copyright	© Copyright © 2018 The Royal Society. All rights reserved. <a href="#">Terms, conditions and policies</a> .

# Pore formation by dimeric Bak and Bax: an unusual pore?

Rachel T. Uren<sup>1,2</sup>, Sweta Iyer<sup>1,2</sup> and Ruth M. Kluck<sup>1,2,3\*</sup>

<sup>1</sup>*Molecular Genetics of Cancer Division, The Walter and Eliza Hall Institute of Medical Research, 1G Royal Parade, Parkville, Victoria, 3052, Australia*

<sup>2</sup>*Department of Medical Biology, The University of Melbourne, 1G Royal Parade, Parkville, Victoria, 3052, Australia*

<sup>3</sup>0000-0002-7101-1925

**Keywords:** apoptosis, Bak, Bax, heterogeneity, membrane bilayer, mitochondrial pore

---

\*Author for correspondence (kluck@wehi.edu.au).

## Summary

Apoptotic cell death via the mitochondrial pathway occurs in all vertebrate cells and requires the formation of pores in the mitochondrial outer membrane. Two Bcl-2 protein family members, Bak and Bax, form these pores during apoptosis and how they do so has been investigated for the last two decades. Many of the conformation changes that occur during their transition to pore-forming proteins have now been delineated. Notably, biochemical, biophysical and structural studies indicate that symmetric homodimers are the basic unit of pore formation. Each dimer contains an extended hydrophobic surface that lies on the outer membrane, and is anchored at either end by a transmembrane domain. Membrane remodelling events such as positive membrane curvature have been reported to accompany apoptotic pore formation, suggesting Bak and Bax form lipidic pores rather than proteinaceous pores. However, it remains unclear how symmetric dimers assemble to porate the membrane. Here we review how clusters of dimers and their lipid-mediated interactions provide a molecular explanation for the heterogeneous assemblies of Bak and Bax observed during apoptosis.

## Introduction

Apoptosis is essential for normal development and tissue homeostasis, and its perturbed regulation contributes to numerous pathological conditions, including cancer and autoimmune and degenerative diseases [1]. Apoptosis is regulated principally by interactions within the Bcl-2 family of proteins, whose members fall into three subclasses (Figure 1a). The eight or more pro-apoptotic BH3-only proteins (e.g. Bid and Bim) act as sensors of specific types of cellular stress, and signal by engaging other family members. The pro-survival proteins (e.g. Bcl-2 and Mcl-1) act by sequestering the pro-apoptotic members. Finally, pro-apoptotic Bak and Bax act as critical effectors of apoptosis, as they are required for mitochondrial permeabilisation in cells and in mice [2, 3]. As illustrated in Figure 1b, upon receiving an apoptotic stimulus, upregulated BH3-only proteins bind to Bak and Bax to induce major conformation changes, resulting in Bak and Bax oligomerisation and subsequent outer membrane permeabilisation. This leads to the release of mitochondrial proteins including cytochrome *c* that in turn triggers caspase-driven cell demolition (reviewed in [4]).

The 3D structures of non-activated Bak and Bax resemble those of the pro-survival proteins, comprising nine  $\alpha$ -helices that form a tight globular bundle (Figure 2). Two important features are a surface hydrophobic groove ( $\alpha 2$ - $\alpha 5$ ) and a buried BH3 domain in  $\alpha 2$  that mediate contact with other family members. A major distinction between Bak and Bax is that Bak is mostly inserted into the mitochondrial outer membrane in healthy cells, whereas Bax is mostly cytosolic and translocates to mitochondria following apoptotic stimuli (Figure 1b). Bax translocation is triggered by binding of BH3-only proteins which releases  $\alpha 9$  from the hydrophobic groove [5]. Once the released  $\alpha 9$  inserts as a transmembrane domain into the mitochondrial outer membrane, Bax has the same topology as non-activated Bak [6].

### Conversion of Bak and Bax into symmetric homodimers with flexible extremities

Bak and Bax undergo major conformation changes as they convert into pore-forming proteins (Figure 2; reviewed in [4]). The changes are triggered by the binding of BH3-only proteins to a hydrophobic surface groove, which generates a cavity underneath both the N- and C-termini [7-9]. Destabilisation allows the protein to unfold as three segments: the  $\alpha 1$ -helix dissociates [10], and the core ( $\alpha 2$ - $\alpha 5$ ) separates from the latch ( $\alpha 6$ - $\alpha 9$ ) [7, 8]. Several newly exposed hydrophobic regions then associate with the mitochondrial outer membrane to lie in-plane (Figure 2, activated Bak monomer)[11, 12]. The core remains largely folded, but within it the newly exposed hydrophobic BH3 domain (in  $\alpha 2$ ) then binds to the hydrophobic groove of another activated Bak or Bax molecule in a reciprocal manner to form symmetric homodimers (Figure 2, BH3:groove dimer). Evidence for symmetric homodimers originated from biochemical studies in mitochondria [13-15] and is supported by X-ray structures of the  $\alpha 2$ - $\alpha 5$  dimers [7, 8] and biophysical studies[12, 16-20].

1  
2  
3 Together these studies support the in-plane model of a Bak dimer (Figure 3a) [11, 21]. The region in  
4 contact with the membrane ( $\alpha 2$ - $\alpha 9$ ) resembles an extended flexible amphipathic polypeptide  
5 anchored at either end with a transmembrane domain - an unusual structure for a pore-forming  
6 protein. Several helices may embed into the outer leaflet of the membrane, encouraged by aromatic  
7 residues on one surface of the  $\alpha 2$ - $\alpha 5$  core dimer (Figure 3b) and on one face of the  $\alpha 6$ - $\alpha 8$  helices  
8 (Figure 3c). At the N-terminus, the first 70 residues become exposed and do not re-engage with either  
9 membrane or protein [10, 21, 22]. Bax dimers also display aromatic residues on one surface (Figure  
10 3b,c) and a similar membrane topology [6, 11, 16, 19], although complete solvent exposure of the N-  
11 terminus has not yet been shown. Thus, Bak and Bax homodimers show several features of  
12 antimicrobial peptides such as human LL-37 and magainin 2 (Figure 3d) that are proposed to form  
13 toroidal pores, rather than of the  $\alpha$ -helical ClyA and actinoporin proteins that form more structured  
14 proteinaceous pores [23, 24].  
15  
16

17  
18  
19  
20  
21  
22  
23  
24 There are few examples of a homodimer as the building block of a pore that might provide insight  
25 into pore formation by Bak and Bax. One such example is the plant defensin NaD1, whose structure  
26 comprises seven antiparallel dimers [25]. However, unlike Bak and Bax, the NaD1 complexes are not  
27 promoted by major conformation change but by binding of the PIP<sub>2</sub> phospholipid, and the oligomers  
28 lack transmembrane domains and flexible membrane-associated regions. Members of the colicin  
29 family of pore-forming proteins may also assemble as multiples of dimers to form small ring-shaped  
30 oligomers (~8 nm in diameter) [24, 26, 27]. Curiously, the first structure of a Bcl-2 protein, and its  
31 similarity to the colicins and diphtheria toxin prompted the idea that, as proposed for those proteins,  
32 Bak and Bax might form channels or pores by inserting a helical hairpin ( $\alpha 5/\alpha 6$ ) through the  
33 membrane [28-30]. Notably, hairpin insertion is not consistent with the in-plane model in which  $\alpha 5$   
34 remains with the  $\alpha 2$ - $\alpha 5$  dimer and  $\alpha 6$  lies in-plane in the outer membrane (Figures 2-4) [11, 12]. A  
35 recent study proposed that Bax  $\alpha 2$ - $\alpha 5$  dimers can progress to  $\alpha 2$ - $\alpha 3$ - $\alpha 4$  dimers after separation of  $\alpha 5$   
36 to allow  $\alpha 5/\alpha 6$  insertion [16]. However, it was not clear whether those dimers were functional [16],  
37 and in other studies separation of Bax  $\alpha 5$  from  $\alpha 4$  was not required for cytochrome *c* release [7].  
38 Even so, further comparison of pore formation by the Bcl-2 and colicin families may prove  
39 informative.  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50

### 51 **Dimer-dimer interactions are not via a single protein-protein interface**

52  
53  
54 It is thought that homodimers of Bak or Bax must then associate to higher order oligomers to porate  
55 the mitochondrial outer membrane. Such oligomers of Bak and of Bax are generated in mitochondria  
56 during apoptosis, as evident by gel filtration, blue native PAGE and linkage studies [31-33]. In  
57 addition, recombinant Bak and Bax form high order oligomers in liposome experiments (table 1).  
58  
59  
60

1 High order oligomers observed biochemically, correlate with the clusters observed in early  
2 microscopy studies (table 1) and that are sometimes apparent at points of mitochondrial fission/fusion  
3 [34]. Higher resolution microscopy shows Bax and Bak complexes of various shapes and sizes in  
4 liposomes and mitochondria, including clusters, rings and arcs (table 1). Thus, there is strong  
5 correlation between high order oligomers and pore formation. Nevertheless, it is yet to be shown that  
6 specifically inhibiting dimer-dimer interaction blocks pore formation.  
7  
8  
9  
10

11  
12 While either Bak or Bax is sufficient to form pores (table 1), the two proteins locate to the same  
13 complexes in apoptotic cells [31, 33], suggesting that mixtures of the two proteins may be able to  
14 generate pores. The mixtures may include heterodimers of Bak and Bax, although heterodimers form  
15 only a minor population compared to homodimers [13, 35]. The low frequency of heterodimers may  
16 be explained by a degree of incompatibility due to the limited sequence similarity of the BH3  
17 domains and grooves of the two proteins. Mixtures may also include homodimers of Bak and of Bax,  
18 as Bax is able to intermingle with pre-formed Bak dimers [21]. If mixtures of Bak and Bax  
19 homodimers can actually generate pores, this would provide further evidence that protein-protein  
20 interactions between dimers are not important for high order oligomers or pore formation.  
21  
22  
23  
24  
25  
26  
27  
28  
29

30 Molecular structures of Bak or Bax as high order oligomers or pore complexes are currently not  
31 available. However, a range of biochemical approaches have been used to examine how activated  
32 Bak and Bax interact to generate pores. Most prominent have been linkage studies showing that  
33 homodimers can associate via interactions at  $\alpha$ -helices 1, 3, 5, 6, and 9 [6, 12, 13, 15, 16, 19, 36-39].  
34 Our initial studies showed that linkage between the  $\alpha$ 6-helices could link dimers of Bak and of Bax  
35 [13, 15], suggesting an  $\alpha$ 6: $\alpha$ 6 interface may drive high order oligomers and pore formation. However,  
36 there was no evidence that mutations in  $\alpha$ 6 could block apoptosis [4, 40], and several groups reported  
37 linkage between additional regions. Thus, some or all of these linkages may be due to collisions  
38 rather than to stable complex formation. Flexibility of the N- and C-termini as depicted in Figure 4a,  
39 may allow linkage between multiple regions, and would also limit interaction between the  $\alpha$ 2- $\alpha$ 5 core  
40 dimers [8, 21]. Based on the linkage pattern throughout the full-length of Bak, we recently proposed  
41 that dimers form disordered clusters during apoptosis (Figure 4a), and this was supported by  
42 mathematical simulation of linkage within the whole population of Bak dimers in the sample [21]. It  
43 is yet to be determined if only a small subpopulation of dimers directly participates in a pore  
44 complex. If so, within this subpopulation the core dimers may adopt an ordered arrangement, e.g.  
45 end-to-end or side-by-side.  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57

## 58 **Formation of lipidic (toroidal) pores**

59  
60

1 Several lines of evidence indicate that Bak and Bax form lipidic rather than proteinaceous pores  
2 (table 1). Amphipathic peptides based on the Bak and Bax  $\alpha 5$  and  $\alpha 6$  helices can permeabilise  
3 membranes (table 1), as do amphipathic antimicrobial peptides that act via forming lipidic pores. As  
4 the Bak and Bax dimers resemble a flexible amphipathic polypeptide (Figure 3a), their shallow  
5 insertion into the outer leaflet [11, 18, 19] may destabilise the lamellar structure of the bilayer to  
6 induce lipidic pores. This mechanism of pore formation may be related to the “carpet” model  
7 proposed for antimicrobial peptides. In the Shai-Matsuzaki-Huang version of the carpet model [41-  
8 43] peptides can disrupt membranes without disintegrating the membranes in a detergent-like manner  
9 [44]. The peptides insert close to the membrane surface to promote a convex curvature of the outer  
10 leaflet. As the peptide concentration increases, membrane defects occur, and in some cases may be  
11 resolved by peptide (or phospholipid) equilibrating across the bilayer. As Bak and Bax are unlikely to  
12 equilibrate across the bilayer due to their size and transmembrane domains, the membrane defects  
13 may progress to pore formation. There is evidence that pore formation is associated with lipid  
14 transbilayer movement [45-47].

25  
26 During pore formation, parts of the dimer may line and stabilise the pore (Figure 4b)[11, 19, 48].  
27 According to the clamp model (Figure 4b, right)[19] the core dimer positions roughly perpendicular  
28 in a circle to line the pore. Several features of this model are attractive. The length of the core dimer  
29 (~4 nm) is the approximate width of the MOM, and the bend observed in the structures (Bak and Bax  
30  $\alpha 2$ - $\alpha 5$  dimers; Figure 3b) may be accommodated by the curved edge of the pore. In this position, the  
31  $\alpha 6$ - $\alpha 8$  latch would disturb the outer and inner leaflets equally, and the core dimer could contribute a  
32 large surface area to stabilise the pore. In addition, the core dimers could pack tightly side-by-side  
33 around the pore. However, one side-by-side orientation of Bak  $\alpha 2$ - $\alpha 5$  dimers observed in a crystal  
34 structure was not supported by linkage studies in mitochondria [8, 21]. Moreover, the clamp model  
35 suggests that the  $\alpha 9$  transmembrane domains become antiparallel within a dimer, presumably after  
36 the charged residues (e.g. RRFFKS in human Bak) at the far C-terminus of one activated molecule  
37 flips through the hydrophobic bilayer. While such flipping may be similar to insertion of the  
38 transmembrane domains of non-activated Bak (and Bax), direct evidence of antiparallel  $\alpha 9$ -helices in  
39 Bax or Bak oligomers is required to support this model.

### 50 51 **Heterogeneity of Bak and Bax complexes and pores**

52 Consistent with forming a lipidic pore, significant heterogeneity is observed in the characteristics of  
53 the Bak and Bax complexes and the actual pores formed by Bak and by Bax (table 1). Differences  
54 may be due to the levels of Bak and Bax (and prosurvival Bcl-2 proteins), lipid composition and  
55 diameter of the mitochondria or liposome, and even the presence or absence of the mitochondrial  
56 inner membrane and matrix. Detectable pores in the membrane were not always evident in the  
57  
58  
59  
60

1 clusters, suggesting that at least some clusters may form upstream or downstream of pore formation.  
2 Notably, Bax pore size in liposomes increased with higher protein concentration and over time [49-  
3 52], and pore size in *Xenopus laevis* mitochondria increased in response to a cytosolic factor that was  
4 more potent in the presence of caspase inhibitors [53]. Thus, both protein and lipid appear able to  
5 enlarge apoptotic pores, a process that would ensure rapid cell death. It will be interesting to  
6 determine the role of pores that are not detectable by microscopy. Can a pore stay small, and what is  
7 the composition of such a "minimal" pore? Are there multiple small pores in a mitochondrion,  
8 perhaps even in a single cluster? Might a single small pore in each mitochondrion be sufficient for  
9 apoptosis? Answers to these questions may help to regulate apoptosis at the step of pore formation,  
10 including reversing the process.  
11  
12  
13  
14  
15  
16  
17  
18  
19

### 20 **Concluding remarks**

21 Increasing evidence indicates that symmetric homodimers of Bak and of Bax form the structural  
22 building block of the apoptotic pore. As there is a strong correlation of higher order oligomers with  
23 pore formation, it is important to understand how symmetric dimers can form these oligomers.  
24 Within each dimer, the N-terminus is solvent-exposed and flexible, implicating the membrane-  
25 associated regions ( $\alpha 2$ - $\alpha 9$ ) in driving pore formation. Flexibility of the latch ( $\alpha 6$ - $\alpha 9$ ) implies several  
26 arrangements of dimers may occur and contribute to the heterogeneity of clusters and pores observed.  
27 Notably, insertion of the core and latch into the outer leaflet may remodel the bilayer to form a small  
28 lipidic pore, which may then grow considerably. Obtaining structures of small apoptotic pores may  
29 yet be possible, and their formation may be the key to understanding how this central step of cell  
30 death might be regulated.  
31  
32  
33  
34  
35  
36  
37  
38  
39

### 40 **Acknowledgments**

41 The authors are grateful for support from NHMRC project grants 637337 and 1008434 and program grant 1016701, the Victorian State  
42 Government Operational Infrastructure Support and the Australian Government NHMRC IRIISS.

### 43 **Authors' Contributions**

44 The authors contributed equally to the planning, writing and illustrating of this article.

### 45 **Competing Interests**

46 The authors declare no competing interests.

### 47 **Funding**

48 Our work is supported by NHMRC grants (637337 and 1016701 to RMK) and by the Victorian State Government Operational  
49 Infrastructure Support and the Australian Government NHMRC IRIISS.  
50  
51  
52

### 53 **References**

- 54 [1] Czabotar, P. E., Lessene, G., Strasser, A. & Adams, J. M. 2014 Control of apoptosis by the BCL-  
55 2 protein family: implications for physiology and therapy. *Nat. Rev. Mol. Cell Biol.* **15**, 49-63.  
56 (DOI:10.1038/nrm3722).  
57 [2] Lindsten, T., Ross, A. J., King, A., Zong, W., Rathmell, J. C., Shiels, H. A., Ulrich, E., Waymire,  
58 K. G., Mahar, P., Frauwirth, K., et al. 2000 The combined functions of proapoptotic Bcl-2 family  
59  
60

- 1 members Bak and Bax are essential for normal development of multiple tissues. *Mol. Cell* **6**, 1389-  
2 1399.
- 3 [3] Wei, M. C., Zong, W. X., Cheng, E. H., Lindsten, T., Panoutsakopoulou, V., Ross, A. J., Roth, K.  
4 A., MacGregor, G. R., Thompson, C. B. & Korsmeyer, S. J. 2001 Proapoptotic BAX and BAK: a  
5 requisite gateway to mitochondrial dysfunction and death. *Science* **292**, 727-730.
- 6 [4] Westphal, D., Kluck, R. M. & Dewson, G. 2014 Building blocks of the apoptotic pore: how Bax  
7 and Bak are activated and oligomerize during apoptosis. *Cell Death Differ.* **21**, 196-205.  
8 (DOI:10.1038/cdd.2013.139).
- 9 [5] Walensky, L. D. & Gavathiotis, E. 2011 BAX unleashed: the biochemical transformation of an  
10 inactive cytosolic monomer into a toxic mitochondrial pore. *Trends Biochem. Sci.* **36**, 642-652.  
11 (DOI:10.1016/j.tibs.2011.08.009).
- 12 [6] Iyer, S., Bell, F., Westphal, D., Anwari, K., Gulbis, J., Smith, B. J., Dewson, G. & Kluck, R. M.  
13 2015 Bak apoptotic pores involve a flexible C-terminal region and juxtaposition of the C-terminal  
14 transmembrane domains. *Cell Death Differ.* (DOI:10.1038/cdd.2015.15).
- 15 [7] Czabotar, P. E., Westphal, D., Dewson, G., Ma, S., Hockings, C., Fairlie, W. D., Lee, E. F., Yao,  
16 S., Robin, A. Y., Smith, B. J., et al. 2013 Bax Crystal Structures Reveal How BH3 Domains Activate  
17 Bax and Nucleate Its Oligomerization to Induce Apoptosis. *Cell* **152**, 519-531.  
18 (DOI:10.1016/j.cell.2012.12.031).
- 19 [8] Brouwer, J. M., Westphal, D., Dewson, G., Robin, A. Y., Uren, R. T., Bartolo, R., Thompson, G.  
20 V., Colman, P. M., Kluck, R. M. & Czabotar, P. E. 2014 Bak Core and Latch Domains Separate  
21 during Activation, and Freed Core Domains Form Symmetric Homodimers. *Mol. Cell* **55**, 938-946.  
22 (DOI:10.1016/j.molcel.2014.07.016).
- 23 [9] Moldoveanu, T., Grace, C. R., Llambi, F., Nourse, A., Fitzgerald, P., Gehring, K., Kriwacki, R.  
24 W. & Green, D. R. 2013 BID-induced structural changes in BAK promote apoptosis. *Nat. Struct.*  
25 *Mol. Biol.* **20**, 589-597. (DOI:10.1038/nsmb.2563).
- 26 [10] Alsop, A. E., Fennell, S. C., Bartolo, R. C., Tan, I. K., Dewson, G. & Kluck, R. M. 2015  
27 Dissociation of Bak alpha1 helix from the core and latch domains is required for apoptosis. *Nat*  
28 *Commun* **6**, 6841. (DOI:10.1038/ncomms7841).
- 29 [11] Westphal, D., Dewson, G., Menard, M., Frederick, P., Iyer, S., Bartolo, R., Gibson, L., Czabotar,  
30 P. E., Smith, B. J., Adams, J. M., et al. 2014 Apoptotic pore formation is associated with in-plane  
31 insertion of Bak or Bax central helices into the mitochondrial outer membrane. *Proc Natl Acad Sci U*  
32 *SA* **111**, E4076-4085. (DOI:10.1073/pnas.1415142111).
- 33 [12] Aluvila, S. M., T. Hustedt, E. Fajer, P. Choe, JY. Oh, KJ. 2014 Organization of the  
34 mitochondrial apoptotic BAK pore: oligomerization of the Bak homodimers. *J. Biol. Chem.* **289**,  
35 2537-2551.
- 36 [13] Dewson, G., Ma, S., Frederick, P., Hockings, C., Tan, I., Kratina, T. & Kluck, R. M. 2012 Bax  
37 dimerizes via a symmetric BH3:groove interface during apoptosis. *Cell Death Differ.* **19**, 661-670.  
38 (DOI:10.1038/cdd.2011.138).
- 39 [14] Dewson, G., Kratina, T., Sim, H. W., Puthalakath, H., Adams, J. M., Colman, P. M. & Kluck, R.  
40 M. 2008 To trigger apoptosis Bak exposes its BH3 domain and homo-dimerizes via BH3:groove  
41 interactions. *Mol. Cell* **30**, 369-380.
- 42 [15] Dewson, G., Kratina, T., Czabotar, P., Day, C. L., Adams, J. M. & Kluck, R. M. 2009 Bak  
43 activation for apoptosis involves oligomerization of dimers via their alpha6 helices. *Mol. Cell* **36**,  
44 696-703. (DOI:S1097-2765(09)00821-1 [pii]  
45 10.1016/j.molcel.2009.11.008).
- 46 [16] Zhang, Z., Subramaniam, S., Kale, J., Liao, C., Huang, B., Brahmabhatt, H., Condon, S. G.,  
47 Lapolla, S. M., Hays, F. A., Ding, J., et al. 2016 BH3-in-groove dimerization initiates and helix 9  
48 dimerization expands Bax pore assembly in membranes. *EMBO J.* **35**, 208-236.  
49 (DOI:10.15252/embj.201591552).
- 50 [17] Subburaj, Y., Cosentino, K., Axmann, M., Pedrueza-Villalmanzo, E., Hermann, E., Bleicken, S.,  
51 Spatz, J. & Garcia-Saez, A. J. 2015 Bax monomers form dimer units in the membrane that further  
52 self-assemble into multiple oligomeric species. *Nat Commun* **6**, 8042. (DOI:10.1038/ncomms9042).
- 53 [18] Oh, K. J., Singh, P., Lee, K., Foss, K., Lee, S., Park, M., Aluvila, S., Kim, R. S., Symersky, J. &  
54 Walters, D. E. 2010 Conformational changes in BAK, a pore-forming proapoptotic Bcl-2 family  
55  
56  
57  
58  
59  
60



- 1 member, upon membrane insertion and direct evidence for the existence of BH3-BH3 contact  
2 interface in BAK homo-oligomers. *J. Biol. Chem.* **285**, 28924-28937. (DOI:M110.135293 [pii]  
3 10.1074/jbc.M110.135293).
- 4 [19] Bleicken, S., Jeschke, G., Stegmüller, C., Salvador-Gallego, R., Garcia-Saez, A. J. &  
5 Bordignon, E. 2014 Structural model of active bax at the membrane. *Mol. Cell* **56**, 496-505.  
6 (DOI:10.1016/j.molcel.2014.09.022).
- 7 [20] Bleicken, S., Classen, M., Padmavathi, P. V., Ishikawa, T., Zeth, K., Steinhoff, H. J. &  
8 Bordignon, E. 2010 Molecular details of Bax activation, oligomerization, and membrane insertion. *J.*  
9 *Biol. Chem.* **285**, 6636-6647. (DOI:M109.081539 [pii]  
10 10.1074/jbc.M109.081539).
- 11 [21] Uren, R. T., O'Hely, M., Iyer, S., Bartolo, R., Brouwer, J. M., Alsop, A. E., Dewson, G. &  
12 Kluck, R. M. 2016 Disordered clusters of Bak dimers rupture mitochondria during apoptosis.  
13 *bioRxiv*. (DOI:10.1101/059899).
- 14 [22] Weber, K., Harper, N., Schwabe, J. & Cohen, G. M. 2013 BIM-mediated membrane insertion of  
15 the BAK pore domain is an essential requirement for apoptosis. *Cell Rep* **5**, 409-420.  
16 (DOI:10.1016/j.celrep.2013.09.010).
- 17 [23] Gilbert, R. J. 2016 Protein-lipid interactions and non-lamellar lipidic structures in membrane  
18 pore formation and membrane fusion. *Biochim. Biophys. Acta* **1858**, 487-499.  
19 (DOI:10.1016/j.bbamem.2015.11.026).
- 20 [24] Dal Peraro, M. & van der Goot, F. G. 2016 Pore-forming toxins: ancient, but never really out of  
21 fashion. *Nat. Rev. Microbiol.* **14**, 77-92. (DOI:10.1038/nrmicro.2015.3).
- 22 [25] Poon, I., Baxter, A. A., Lay, F. T., Mills, G. D., Adda, C. G., Payne, J. A., Phan, T. K., Ryan, G.  
23 F., White, J. A., Veneer, P. K., et al. 2014 Phosphoinositide-mediated oligomerization of a defensin  
24 induces cell lysis. *Elife* **3**, e01808. (DOI:10.7554/eLife.01808).
- 25 [26] Greig, S. L., Radjainia, M. & Mitra, A. K. 2009 Oligomeric structure of colicin ia channel in  
26 lipid bilayer membranes. *J. Biol. Chem.* **284**, 16126-16134. (DOI:10.1074/jbc.M900292200).
- 27 [27] Dunkel, S., Pulagam, L. P., Steinhoff, H. J. & Klare, J. P. 2015 In vivo EPR on spin labeled  
28 colicin A reveals an oligomeric assembly of the pore-forming domain in E. coli membranes. *Phys*  
29 *Chem Chem Phys* **17**, 4875-4878. (DOI:10.1039/e4cp05638h).
- 30 [28] Muchmore, S. W., Sattler, M., Liang, H., Meadows, R. P., Harlan, J. E., Yoon, H. S.,  
31 Nettlesheim, D., Chang, B. S., Thompson, C. B., Wong, S.-L., et al. 1996 X-ray and NMR structure of  
32 human Bcl-x<sub>L</sub>, an inhibitor of programmed cell death. *Nature* **381**, 335-341.
- 33 [29] Antignani, A. & Youle, R. J. 2006 How do Bax and Bak lead to permeabilization of the outer  
34 mitochondrial membrane? *Curr. Opin. Cell Biol.* **18**, 685-689. (DOI:S0955-0674(06)00157-8 [pii]  
35 10.1016/j.ceb.2006.10.004).
- 36 [30] Annis, M. G., Soucie, E. L., Dlugosz, P. J., Cruz-Aguado, J. A., Penn, L. Z., Leber, B. &  
37 Andrews, D. W. 2005 Bax forms multispanning monomers that oligomerize to permeabilize  
38 membranes during apoptosis. *EMBO J.* **24**, 2096-2103.
- 39 [31] Zhou, L. & Chang, D. C. 2008 Dynamics and structure of the Bax-Bak complex responsible for  
40 releasing mitochondrial proteins during apoptosis. *J. Cell Sci.* **121**, 2186-2196.
- 41 [32] Wei, M. C., Lindsten, T., Mootha, V. K., Weiler, S., Gross, A., Ashiya, M., Thompson, C. B. &  
42 Korsmeyer, S. J. 2000 tBID, a membrane-targeted death ligand, oligomerizes BAK to release  
43 cytochrome c. *Genes Dev.* **14**, 2060-2071.
- 44 [33] Nechushtan, A., Smith, C. L., Lamensdorf, I., Yoon, S. H. & Youle, R. J. 2001 Bax and Bak  
45 coalesce into novel mitochondria-associated clusters during apoptosis. *J. Cell Biol.* **153**, 1265-1276.
- 46 [34] Karbowski, M., Lee, Y. J., Gaume, B., Jeong, S. Y., Frank, S., Nechushtan, A., Santel, A.,  
47 Fuller, M., Smith, C. L. & Youle, R. J. 2002 Spatial and temporal association of Bax with  
48 mitochondrial fission sites, Drp1, and Mfn2 during apoptosis. *J. Cell Biol.* **159**, 931-938.  
49 (DOI:10.1083/jcb.200209124).
- 50 [35] Mikhailov, V., Mikhailova, M., Degenhardt, K., Venkatachalam, M. A., White, E. & Saikumar,  
51 P. 2003 Association of Bax and Bak homo-oligomers in mitochondria. Bax requirement for Bak  
52 reorganization and cytochrome c release. *J. Biol. Chem.* **278**, 5367-5376.  
53 (DOI:10.1074/jbc.M203392200).
- 54  
55  
56  
57  
58  
59  
60

- [36] Zhang, Z., Zhu, W., Lapolla, S. M., Miao, Y., Shao, Y., Falcone, M., Boreham, D., McFarlane, N., Ding, J., Johnson, A. E., et al. 2010 Bax forms an oligomer via separate, yet interdependent, surfaces. *J. Biol. Chem.* **285**, 17614-17627. (DOI:10.1074/jbc.M110.113456 [pii] 10.1074/jbc.M110.113456).
- [37] Pang, Y. P., Dai, H., Smith, A., Meng, X. W., Schneider, P. A. & Kaufmann, S. H. 2012 Bak Conformational Changes Induced by Ligand Binding: Insight into BH3 Domain Binding and Bak Homo-Oligomerization. *Sci Rep* **2**, 257. (DOI:10.1038/srep00257).
- [38] Ma, S., Hockings, C., Anwari, K., Kratina, T., Fennell, S., Lazarou, M., Ryan, M. T., Kluck, R. M. & Dewson, G. 2013 Assembly of the Bak apoptotic pore: a critical role for the Bak protein alpha6 helix in the multimerization of homodimers during apoptosis. *J. Biol. Chem.* **288**, 26027-26038. (DOI:10.1074/jbc.M113.490094).
- [39] Gahl, R. F., He, Y., Yu, S. & Tjandra, N. 2014 Conformational rearrangements in the pro-apoptotic protein, Bax, as it inserts into mitochondria: a cellular death switch. *J. Biol. Chem.* **289**, 32871-32882. (DOI:10.1074/jbc.M114.593897).
- [40] Westphal, D., Dewson, G., Czabotar, P. E. & Kluck, R. M. 2011 Molecular biology of Bax and Bak activation and action. *Biochim. Biophys. Acta* **1813**, 521-531. (DOI:S0167-4889(10)00336-8 [pii] 10.1016/j.bbamcr.2010.12.019).
- [41] Gazit, E., Miller, I. R., Biggin, P. C., Sansom, M. S. & Shai, Y. 1996 Structure and orientation of the mammalian antibacterial peptide cecropin P1 within phospholipid membranes. *J. Mol. Biol.* **258**, 860-870. (DOI:10.1006/jmbi.1996.0293).
- [42] Ludtke, S. J., He, K., Heller, W. T., Harroun, T. A., Yang, L. & Huang, H. W. 1996 Membrane pores induced by magainin. *Biochemistry* **35**, 13723-13728. (DOI:10.1021/bi9620621).
- [43] Matsuzaki, K., Murase, O., Tokuda, H., Funakoshi, S., Fujii, N. & Miyajima, K. 1994 Orientational and aggregational states of magainin 2 in phospholipid bilayers. *Biochemistry* **33**, 3342-3349.
- [44] Lee, M. T., Hung, W. C., Chen, F. Y. & Huang, H. W. 2008 Mechanism and kinetics of pore formation in membranes by water-soluble amphipathic peptides. *Proc Natl Acad Sci U S A* **105**, 5087-5092. (DOI:10.1073/pnas.0710625105).
- [45] Epand, R. F., Martinou, J. C., Montessuit, S. & Epand, R. M. 2003 Transbilayer lipid diffusion promoted by Bax: implications for apoptosis. *Biochemistry* **42**, 14576-14582. (DOI:10.1021/bi035348w).
- [46] Terrones, O., Antonsson, B., Yamaguchi, H., Wang, H. G., Liu, J., Lee, R. M., Herrmann, A. & Basanez, G. 2004 Lipidic pore formation by the concerted action of proapoptotic BAX and tBID. *J Biol Chem* **279**, 30081-30091.
- [47] Garcia-Saez, A. J., Coraiola, M., Serra, M. D., Mingarro, I., Muller, P. & Salgado, J. 2006 Peptides corresponding to helices 5 and 6 of Bax can independently form large lipid pores. *FEBS J.* **273**, 971-981. (DOI:EJB5123 [pii] 10.1111/j.1742-4658.2006.05123.x).
- [48] Mandal, T., Shin, S., Aluvila, S., Chen, H. C., Grieve, C., Choe, J. Y., Cheng, E. H., Hustedt, E. J. & Oh, K. J. 2016 Assembly of Bak homodimers into higher order homooligomers in the mitochondrial apoptotic pore. *Sci Rep* **6**, 30763. (DOI:10.1038/srep30763).
- [49] Korsmeyer, S. J., Wei, M. C., Saito, M., Weiler, S., Oh, K. J. & Schlesinger, P. H. 2000 Pro-apoptotic cascade activates BID, which oligomerizes BAK or BAX into pores that result in the release of cytochrome *c*. *Cell Death Differ.* **7**, 1166-1173.
- [50] Saito, M., Korsmeyer, S. J. & Schlesinger, P. H. 2000 BAX-dependent transport of cytochrome *c* reconstituted in pure liposomes. *Nat. Cell Biol.* **2**, 553-555. (DOI:10.1038/35019596).
- [51] Bleicken, S., Landeta, O., Landajuela, A., Basanez, G. & Garcia-Saez, A. J. 2013 Proapoptotic Bax and Bak proteins form stable protein-permeable pores of tunable size. *J. Biol. Chem.* **288**, 33241-33252. (DOI:10.1074/jbc.M113.512087).
- [52] Gillies, L. A., Du, H., Peters, B., Knudson, C. M., Newmeyer, D. D. & Kuwana, T. 2015 Visual and functional demonstration of growing Bax-induced pores in mitochondrial outer membranes. *Mol Biol Cell* **26**, 339-349. (DOI:10.1091/mbc.E13-11-0638).

- [53] Kluck, R. M., Esposti, M. D., Perkins, G., Renken, C., Kuwana, T., Bossy-Wetzel, E., Goldberg, M., Allen, T., Barber, M. J., Green, D. R., et al. 1999 The pro-apoptotic proteins, Bid and Bax, cause a limited permeabilization of the mitochondrial outer membrane that is enhanced by cytosol. *J. Cell Biol.* **147**, 809-822.
- [54] Qian, S., Wang, W., Yang, L. & Huang, H. W. 2008 Structure of transmembrane pore induced by Bax-derived peptide: evidence for lipidic pores. *Proc Natl Acad Sci U S A* **105**, 17379-17383. (DOI:0807764105 [pii] 10.1073/pnas.0807764105).
- [55] Garcia-Saez, A. J., Coraiola, M., Dalla Serra, M., Mingarro, I., Menestrina, G. & Salgado, J. 2005 Peptides derived from apoptotic Bax and Bid reproduce the poration activity of the parent full-length proteins. *Biophys. J.* **88**, 3976-3990. (DOI:S0006-3495(05)73449-4 [pii] 10.1529/biophysj.104.058008).
- [56] Garcia-Saez, A. J., Chiantia, S., Salgado, J. & Schwille, P. 2007 Pore formation by a Bax-derived peptide: effect on the line tension of the membrane probed by AFM. *Biophys. J.* **93**, 103-112. (DOI:10.1529/biophysj.106.100370).
- [57] Schlesinger, P. H., Gross, A., Yin, X. M., Yamamoto, K., Saito, M., Waksman, G. & Korsmeyer, S. J. 1997 Comparison of the ion channel characteristics of proapoptotic BAX and antiapoptotic BCL-2. *Proc Natl Acad Sci U S A* **94**, 11357-11362.
- [58] Antonsson, B., Conti, F., Ciavatta, A., Montessuit, S., Lewis, S., Martinou, I., Bernasconi, L., Bernard, A., Mermod, J. J., Mazzei, G., et al. 1997 Inhibition of Bax channel-forming activity by Bcl-2. *Science* **277**, 370-372.
- [59] Landeta, O., Landajuena, A., Gil, D., Taneva, S., Di Primo, C., Sot, B., Valle, M., Frolov, V. A. & Basanez, G. 2011 Reconstitution of proapoptotic BAK function in liposomes reveals a dual role for mitochondrial lipids in the BAK-driven membrane permeabilization process. *J. Biol. Chem.* **286**, 8213-8230. (DOI:10.1074/jbc.M110.165852).
- [60] Basanez, G., Sharpe, J. C., Galanis, J., Brandt, T. B., Hardwick, J. M. & Zimmerberg, J. 2002 Bax-type apoptotic proteins porate pure lipid bilayers through a mechanism sensitive to intrinsic monolayer curvature. *J. Biol. Chem.* **277**, 49360-49365. (DOI:10.1074/jbc.M206069200 M206069200 [pii]).
- [61] Basanez, G., Nechushtan, A., Drozhinin, O., Chanturiya, A., Choe, E., Tutt, S., Wood, K. A., Hsu, Y., Zimmerberg, J. & Youle, R. J. 1999 Bax, but not Bcl-xL, decreases the lifetime of planar phospholipid bilayer membranes at subnanomolar concentrations. *Proc Natl Acad Sci U S A* **96**, 5492-5497.
- [62] Epand, R. F., Martinou, J. C., Montessuit, S., Epand, R. M. & Yip, C. M. 2002 Direct evidence for membrane pore formation by the apoptotic protein Bax. *Biochem. Biophys. Res. Commun.* **298**, 744-749.
- [63] Kuwana, T., Mackey, M. R., Perkins, G., Ellisman, M. H., Latterich, M., Schneider, R., Green, D. R. & Newmeyer, D. D. 2002 Bid, Bax, and lipids cooperate to form supramolecular openings in the outer mitochondrial membrane. *Cell* **111**, 331-342.
- [64] Bleicken, S., Wagner, C. & Garcia-Saez, A. J. 2013 Mechanistic differences in the membrane activity of Bax and Bcl-xL correlate with their opposing roles in apoptosis. *Biophys. J.* **104**, 421-431. (DOI:10.1016/j.bpj.2012.12.010).
- [65] Bleicken, S., Hofhaus, G., Ugarte-Urbe, B., Schroder, R. & Garcia-Saez, A. J. 2016 cBid, Bax and Bcl-xL exhibit opposite membrane remodeling activities. *Cell Death Dis* **7**, e2121. (DOI:10.1038/cddis.2016.34).
- [66] Xu, X. P., Zhai, D., Kim, E., Swift, M., Reed, J. C., Volkman, N. & Hanein, D. 2013 Three-dimensional structure of Bax-mediated pores in membrane bilayers. *Cell Death Dis* **4**, e683. (DOI:10.1038/cddis.2013.210).
- [67] Schafer, B., Quispe, J., Choudhary, V., Chipuk, J. E., Ajero, T. G., Du, H., Schneider, R. & Kuwana, T. 2009 Mitochondrial outer membrane proteins assist Bid in Bax-mediated lipidic pore formation. *Mol Biol Cell* **20**, 2276-2285. (DOI:10.1091/mbc.E08-10-1056).
- [68] Kuwana, T., Olson, N. H., Kiosses, W. B., Peters, B. & Newmeyer, D. D. 2016 Pro-apoptotic Bax molecules densely populate the edges of membrane pores. *Sci Rep* **6**, 27299. (DOI:10.1038/srep27299).

- 1 [69] Salvador-Gallego, R., Mund, M., Cosentino, K., Schneider, J., Unsay, J., Schraermeyer, U.,  
2 Engelhardt, J., Ries, J. & Garcia-Saez, A. J. 2016 Bax assembly into rings and arcs in apoptotic  
3 mitochondria is linked to membrane pores. *EMBO J.* **35**, 389-401. (DOI:10.15252/embj.201593384).  
4 [70] Nasu, Y., Benke, A., Arakawa, S., Yoshida, G. J., Kawamura, G., Manley, S., Shimizu, S. &  
5 Ozawa, T. 2016 In Situ Characterization of Bak Clusters Responsible for Cell Death Using Single  
6 Molecule Localization Microscopy. *Sci Rep* **6**, 27505. (DOI:10.1038/srep27505).  
7 [71] Grosse, L., Wurm, C. A., Bruser, C., Neumann, D., Jans, D. C. & Jakobs, S. 2016 Bax  
8 assembles into large ring-like structures remodeling the mitochondrial outer membrane in apoptosis.  
9 *EMBO J.* **35**, 402-413. (DOI:10.15252/embj.201592789).  
10 [72] Kvensakul, M., Yang, H., Fairlie, W. D., Czabotar, P. E., Fischer, S. F., Perugini, M. A., Huang,  
11 D. C. & Colman, P. M. 2008 Vaccinia virus anti-apoptotic F1L is a novel Bcl-2-like domain-  
12 swapped dimer that binds a highly selective subset of BH3-containing death ligands. *Cell Death*  
13 *Differ.* **15**, 1564-1571.  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

For Review  
Author Manuscript

**Table 1.** Heterogeneity of Bak and Bax complexes and pores.

	Bak/Bax	Membrane type	Resolution Method	Characteristics of complexes and pores, including effect on membranes	References
<b>Peptides</b>	Bax $\alpha 5$ , $\alpha 6$	LUV Planar bilayer	AFM CD spectroscopy Confocal X-ray diffraction	Toroidal pores of diameter $\sim 5.8$ nm Decreased membrane line tension at pore rim Lipids with positive intrinsic curvature enhance pore formation Lipid transbilayer redistribution activity for Bax $\alpha 5$	Garcia-Saez 2005, 2006, 2007 Qian 2008 [47, 54-56]
	Bax $\Delta C$ Bax	Liposome Planar bilayer	Patch clamping Fl-dextran release	Bax forms pH dependent ion-conduction channels/pores Maximum of 4 molecules, pore diameter $\sim 2.2$ nm	Antonsson 1997 Schlesinger 1997 Saito 2000 [50, 57, 58]
<b>Recombinant proteins</b>	Bax $\Delta C$ IVT-Bax Bax (bovine)	LUV Liposome Planar bilayer	Voltage clamp Membrane lifetime measurements	Decrease in membrane lifetime and linear tension Increase in positive monolayer curvature stress Lipid transbilayer redistribution	Basanez 1999 Basanez 2002 Terrones 2004 Landeta 2011 [46, 59-61]
	Bax Bax $\Delta C$	LUV	AFM	Toroidal pores of diameter $\sim 100$ -300 nm Small ring-like structures on bilayer surface Structures contain clusters of $\sim 22$ Bax monomers Lipid transbilayer redistribution	Epand 2002 Epand 2003 [45, 62]
	Bax OG-Bax	Liposome OMV ( <i>Xenopus</i> )	TEM Dextran release	Supramolecular lipid pores Allowed release of $\sim 2000$ kDa dextran	Kuwana 2002 [63]
	Bax Bak $\Delta C$	LUV GUV Mitochondria	Cryo-EM FCS	Time and protein concentration dependent lipid pores Pores of diameter $\sim 3$ -140 nm Induction/stabilisation of curved membrane structures Reduction in vesicle size due to budding, fission, tethering	Bleicken 2013a, 2013b, 2016 [51, 64, 65]
	Bax	Bilayer nanodiscs	Cryo-EM	Active Bax monomers form pores of diameter $\sim 3.5$ nm	Xu 2013 [66]
	Bax Nanogold labelled Bax	Liposome LUV OMV ( <i>Xenopus</i> )	TEM Cryo-EM	Pore-like openings with diameter $\sim 25$ -100 nm Growing pores in the range of $\sim 100$ -300 nm Solitary dynamic pores with negative curvature at edges Complete rings of Bax exclusively associated with pore rims	Schafer 2009 Gillies 2015 Kuwana 2016 [52, 67, 68]
	Bax	LUV Planar bilayer	AFM TIRF	Multiple oligomer species of dimers Round heterogeneous pores of diameter $\sim 24$ -176 nm Bax along pore rim	Subburaj 2015 Salvador-Gallego 2016 [17, 69]
<b>Cells</b>	GFP-Bax	HeLa Cos-7	TEM Confocal	Large clusters containing 1000-20,000 Bax molecules	Nechushtan 2001 [33]
	CFP-Bax	HeLa	Quantitative fluorescence imaging	Bax complexes with $\sim 150$ -1000 molecules per cluster 620 nm average cluster size	Zhou 2008 [31]
	mEos3-Bak	MEF	PALM	Heterogeneous Bak clusters of diameter $\sim 70$ -600 nm Each cluster contains $\sim 20$ -2000 Bax molecules No pores evident in clusters (using resolution of 20 nm)	Nasu 2016 [70]
	Endog. Bax	HeLa U2OS HT1080 SH-SY5Y CV-1	STED	Large compact clusters Rings up to 400 nm diameter	Grosse 2016 [71]
	GFP-Bax	HeLa HCT-116	TEM STED SMLM	Heterogenous distribution of: clusters rings (35 nm diameter) arcs (100-500 nm diameter) lines	Salvador-Gallego 2016 [69]

AFM, Atomic Force Microscopy; CD, Circular Dichroism; Cryo-EM, Cryo-Electron Microscopy;  $\Delta C$ , C-terminally truncated; FCS, Fluorescence Correlation Spectroscopy; GUV, Giant Unilamellar Vesicles; IVT, In Vitro Translated; LUV, Large Unilamellar Vesicles; OG, Octyl Glucoside; OMV, Outer Membrane Vesicles; PALM, Photo-activated Localisation Microscopy; TEM, Transmission Electron Microscopy; TIRF, Total Internal Reflection Fluorescence Microscopy; SMLM, Single Molecule Localisation Microscopy; STED, Stimulated Emission Depletion microscopy.

**Figure 1.**

Bcl-2 proteins regulate the mitochondrial pathway of apoptotic cell death. (a) Three subfamilies of Bcl-2 proteins. (b) Bak and Bax activation by the BH3-only proteins is followed by their oligomerisation in the mitochondrial outer membrane to release cytochrome *c* and induce apoptosis.

**Figure 2.**

Bak activation and conformation change results in symmetric homodimers. A schematic showing that Bak unfolds by the N-terminus ( $\alpha 1$ , blue) and the C-terminal latch ( $\alpha 6$ - $\alpha 8$ , magenta) separating from the  $\alpha 2$ - $\alpha 5$  core (orange, red). Hydrophobic regions of the core and latch then collapse onto the membrane, while the exposed BH3 domain (in  $\alpha 2$ ) binds to the hydrophobic groove in another activated Bak molecule. Reciprocal BH3:groove binding results in symmetric homodimers. The indicated crystal structures demonstrate the major conformation changes involved. Equivalent changes are observed for Bax.

**Figure 3.**

Membrane topology of the Bak dimer. (a) The in-plane model of the Bak dimer. The N-terminal regions become solvent-exposed while the remainder of the Bak dimer resembles a flexible extended amphipathic polypeptide that lies in-plane with the membrane, anchored at either end by transmembrane domains. Note that  $\alpha 1$  may unfold after it dissociates, decreasing the hydrophobicity of the BH4 structural motif (VFrsYV) therein [10, 72]. Images were assembled in PyMol using the structures of Bak (2IMT) and the Bak dimer (4U2V), and represented as cartoon and mesh. Colour coding as in Figure 2. (b) Aromatic residues are concentrated on the bent surface of the Bak and Bax  $\alpha 2$ - $\alpha 5$  core dimers. (c) Aromatic residues can position on one edge of the flexible  $\alpha 6$ - $\alpha 8$  latch. (d) Examples of antimicrobial peptides thought to form lipidic pores, with aromatic residues indicated.

**Figure 4.**

Possible mechanisms involved in lipidic pore formation and stabilisation by homodimers of Bak or Bax. (a) Schematic of Bak dimers forming disordered clusters on the mitochondrial outer membrane, encouraged by flexibility of the  $\alpha 6$ - $\alpha 8$  latch. Note that end-to-end or side-by-side contact between the core regions is possible. Images were assembled as in Fig. 3a. (b) Parts of the dimer may line a lipidic pore. The flexible amphipathic latch may slide into a nascent pore to partially line and stabilise the pore (left; in-plane model [4]). The amphipathic core dimer ( $\alpha 2$ - $\alpha 5$ ) may also line the pore generating anti-parallel  $\alpha 9$ -helices (right; clamp model [19]).

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60



Figure 1

Figure 1

297x420mm (300 x 300 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

Figure 2




Figure 2

297x420mm (300 x 300 DPI)






Figure 3

297x420mm (300 x 300 DPI)

1  
2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60

A Disordered clusters of Bak dimers  
(top view, NT removed)





Figure 4

B Stabilisation of lipidic pore



Author Manuscript

Figure 4

297x420mm (300 x 300 DPI)